

Neuromechanical adaptations in joint stiffness when hopping with a SL-AFO


Presentation by Ronald A. Roiz
Research conducted by: Ronald A. Roiz, Arick Auyang,
Dr. Y.H. Chang
Georgia Institute of Technology
GIT IRB # H04189
04.11.2007

Introduction


- The human body is modeled to behave like a spring-mass system during bouncing gaits such as running and hopping.^{1-4,21, 23}

- The **leg's stiffness** influences:^{2,3,5-7}

- peak ground reaction force
- ground contact time
- center of mass displacement
- stride frequency
- “stiffness is the primary control variable in motor behavior.”⁹




Ferris et al, 2006



Significance


- Basic Science Question:
 - Understand how local control variables are adjusted to maintain stable global tasks.^{11, 19-20}
 - during hopping leg stiffness (global task) is modulated at the ankle (local variable) with different perturbations:^{5-6,8}



Significance

- Basic Science Question:

- Understand how local control variables are adjusted to maintain stable global tasks. ^{11, 19-20}
- during hopping leg stiffness (global task) is modulated at the ankle (local variable) with different perturbations:^{5-6,8}
 - an elastic element in series



[Ferris et al, 1997]

Significance

- Basic Science Question:

- Understand how local control variables are adjusted to maintain stable global tasks. ^{11, 19-20}
- during hopping leg stiffness (global task) is modulated at the ankle (local variable) with different perturbations:^{5-6,8}
- an elastic element in series



Significance

- Basic Science Question:


- Understand how local control variables are adjusted to maintain stable global tasks. ^{11, 19-20}
- during hopping leg stiffness (global task) is modulated at the ankle (local variable) with different perturbations:^{5-6,8}
 - an elastic element in parallel ⁵



Hypothesis

Hypothesis A: When a spring is added in parallel to the leg to create a *plantar-flexor* torque, the effective leg stiffness will stay the same, but the biological ankle stiffness will *decrease*.

Hypothesis B: When a spring is added in parallel to the leg to create a *dorsi-flexor* torque, the effective leg stiffness will stay the same, but the biological ankle stiffness will *increase*



Research Design & Methods

■ Subjects[‡] :

- Inclusion Criteria: no major muscular or skeletal injuries, or neurologic conditions
- 10 Total: 6 Male, 4 Female
- Avg. Weight: 61.34 Kg (11.16) Avg. Age: 25.4 y/o (2.46)

■ Protocol:

- 3 conditions^{*‡}: AFO, AFO w/ PFA, AFO w/PFR
- 3 frequencies for each condition^{**‡}: 2.2 Hz, 2.4 Hz, and 2.8 Hz
- 3 trials for each condition-frequency combination
- 10 sec of hopping on Right Leg for each trial when frequency is reached
- ~30 hops were analyzed per condition

* 3 trials taken at preferred frequency with no AFO

** 3 trials taken at 2.2Hz with no AFO

[‡]INDEPENDENT VARIABLES

■ Instrumentation used: Vicon Motion Capture system, AMTI Force Plate, Konigsberg 8-channel bi-polar EMG, and an Omega Force Transducer

■ Statistical Methods

- Three-Way AVOVA: Repeated Measures
- Post-Hoc: Bonferroni

General Procedures:



Data Analysis

Dependent Variables:

Leg stiffness = Linear Regression ($\Delta \text{vGRF} / \Delta \text{COM}$)

- Units: N / m
- vGRF (from Force plate)
- $\text{COM} = \iint \text{vGRF}/m \, dt^2$

Total Joint stiffness = Linear Regression ($\Delta M / \Delta \theta$)

- Units: Nm / rad
- M (inverse kinematics)
- $\Delta \theta$ (from Vicon)


Orthosis Ankle Stiffness = Linear Regression ($\Delta F_{FT} \times d_{sp} / \Delta \theta$)

- Units: Nm / rad
- F_{FT} (force transducer)
- d_{sp} (geometric model of spring moment arm)


Biological Ankle Stiffness = Total Ankle Stiffness – Orthosis Ankle Stiffness

- Units: Nm / rad

Normalized Leg Stiffness (N = 10)



Normalized Total Ankle Stiffness (N=10)



Study Limitations

- AFO may not be universally designed for all subjects
- Spring length predetermined for all subjects, rather than for each person
- Spring stiffness predetermined for all subjects rather than for each person
- Inclusion criteria for data was: Hopping $f \pm 3\%$; $f \propto k^2$ which implies k may vary $\pm 9\%$

Discussion

- Leg stiffness was maintained within a frequency, though it is not mechanically necessary to do so.^{3,7}
- Therefore it is shown that the global task (leg stiffness) is maintained under new perturbation paradigm; i.e., with a dorsi-flexor torque.
- To maintain an invariant effective leg stiffness with this perturbation requires that compensation is occurring at one or more of the joints – though which one it is has yet to be deciphered since it does not appear to be entirely at the ankle.

Application

- Further insight into a rehabilitative modality achieved
 - With variable assistance/resistance we can work toward therapeutic AFOs
- More sophisticated rehabilitation devices could be made effectively. ¹⁴⁻¹⁸
 - Compliance of device (e.g. a PLS-AFO or GRF-AFO) could directly impact the effective stiffness of the joint it is being applied to.



Acknowledgements

- Special thanks to:
 - Dr. Y.H. Chang
 - Arick Auyang
 - Rob A. MacDonald
 - Dr. T. Snow
 - Jasper Yen
 - Heather Brant

and of course my subjects,
who spent 3 hrs. of their
lives with me – a feat in of
itself



References


1. Farley CT, Blickhan R, Saito J, Taylor CR. Hopping frequency in humans: a test of how springs set stride frequency in bouncing gaits. *J Appl Physiol*. Dec 1991;71(6):2127-2132.
2. Farley CT, Houdijk HH, Van Strien C, Louie M. Mechanism of leg stiffness adjustment for hopping on surfaces of different stiffnesses. *J Appl Physiol*. Sep 1998;85(3):1044-1055.
3. Farley CT, Morgenroth DC. Leg stiffness primarily depends on ankle stiffness during human hopping. *J Biomech*. Mar 1999;32(3):267-273.
4. Ferris DP, Liang K, Farley CT. Runners adjust leg stiffness for their first step on a new running surface. *J Biomech*. Aug 1999;32(8):787-794.
5. Ferris DP, Bohra ZA, Lukos JR, Kinnaird CR. Neuromechanical adaptation to hopping with an elastic ankle-foot orthosis. *J Appl Physiol*. Jan 2006;100(1):163-170.
6. Moritz CT, Greene SM, Farley CT. Neuromuscular changes for hopping on a range of damped surfaces. *J Appl Physiol*. May 2004;96(5):1996-2004.
7. Ferris DP, Farley CT. Interaction of leg stiffness and surfaces stiffness during human hopping. *J Appl Physiol*. Jan 1997;82(1):15-22; discussion 13-14.
8. Moritz CT, Farley CT. Human hopping on damped surfaces: strategies for adjusting leg mechanics. *Proc Biol Sci*. Aug 22 2003;270(1525):1741-1746.
9. Granata KP, Wilson SE, Massimini AK, Gabriel R. Active stiffness of the ankle in response to inertial and elastic loads. *J Electromyogr Kinesiol*. Oct 2004;14(5):599-609.
10. Houk JC. Regulation of stiffness by skeletomotor reflexes. *Annu Rev Physiol*. 1979;41:99-114.
11. Lacquaniti F, Licata F, Soechting JF. The mechanical behavior of the human forearm in response to transient perturbations. *Biol Cybern*. 1982;44(1):35-46.
12. Siegler EL, Beck LH. Stiffness: a pathophysiologic approach to diagnosis and treatment. *J Gen Intern Med*. Nov-Dec 1989;4(6):533-540.

References

13. Siegler S, Moskowitz GD, Freedman W. Passive and active components of the internal moment developed about the ankle joint during human ambulation. *J Biomech.* 1984;17(9):647-652.
14. Sawicki GS, Domingo A, Ferris DP. The effects of powered ankle-foot orthoses on joint kinematics and muscle activation during walking in individuals with incomplete spinal cord injury. *J Neuroengineering Rehabil.* 2006;3:3.
15. Ferris DP, Gordon KE, Sawicki GS, Peethambaran A. An improved powered ankle-foot orthosis using proportional myoelectric control. *Gait Posture.* Jun 2006;23(4):425-428.
16. Gordon KE, Sawicki GS, Ferris DP. Mechanical performance of artificial pneumatic muscles to power an ankle-foot orthosis. *J Biomech.* 2006;39(10):1832-1841.
17. Bartonek A, Eriksson M, Gutierrez-Farewik EM. A new carbon fibre spring orthosis for children with plantarflexor weakness. *Gait Posture.* Sep 6 2006.
18. Hollander KW, Ilg R, Sugar TG, Herring D. An efficient robotic tendon for gait assistance. *J Biomech Eng.* Oct 2006;128(5):788-791.
19. Gard SA, Childress DS. The influence of stance-phase knee flexion on the vertical displacement of the trunk during normal walking. *Arch Phys Med Rehabil.* Jan 1999;80(1):26-32.
20. Gard SA, Childress DS. What determines the vertical displacement of the body during normal walking? *JPO.* 2001;13(3):64.
21. Blickhan R. The spring-mass model for running and hopping. *J Biomech.* 1989;22(11-12):1217-1227.
22. Cavagna GA. Force platforms as ergometers. *J Appl Physiol.* Jul 1975;39(1):174-179.
23. McMahon TA, Cheng GC. The mechanics of running: how does stiffness couple with speed? *J Biomech.* 1990;23 Suppl 1:65-78.
24. Mehrholz J, Rutte K, Pohl M. Jump training is feasible for nearly ambulatory patients after stroke. *Clin Rehabil.* May 2006;20(5):406-412.

Use of EMG

- Physiological Check:
 - “joint stiffness” varies with muscle activation⁵
 - variations to be expected with the main flexors/extensors surrounding the ankle and knee




RESISTIVE TORQUE

ASSISTIVE TORQUE

Biological Ankle Stiffness (Nm/rad)

Applied Torque (Nm/rad)



SL-AFO

